

CHAMOT

Recombinant Human IL-5

CM007-5HP
CM007-20HP
CM007-100HP
CM007-500HP
CM007-1000HP



CHAMOT

喬默生物技術(上海)有限公司
CHAMOT BIOTECHNOLOGY CO., LTD.

CONTENT

1 产品简介

2 产品组成

3 产品储存


4 产品使用

Recombinant Human IL-5 (Interleukin-5)

编号:	CM007-5HP CM007-20HP CM007-100HP CM007-500HP CM007-1000HP	规格:	5 µg 20 µg 100 µg 500 µg 1 mg
类别:	重组蛋白	应用:	Functional Assay

产品简介

描述:	Interleukin 5 (IL5) is an interleukin produced by type-2 T helper cells and mast cells. Through binding to the interleukin-5 receptor, interleukin 5 stimulates B cell growth and increases immunoglobulin secretion - primarily IgA. It is also a key mediator in eosinophil activation.
来源:	<i>Escherichia coli</i>
纯度:	>98% as determined by SDS-PAGE. Ni-NTA chromatography
生物学活性:	Measure by its ability to induce TF-1 cells proliferation. The ED ₅₀ for this effect is < 0.3 ng/mL.
内毒素检测:	< 0.1 EU per 1 µg of the protein by the LAL method.
蛋白序列:	MIPTSEPTSAIVKETLALLSTHRTLLIAN ETLRIPVPVHKNHQLCTEEIFQGIGTLE SQTVQGGTVERLFKNLSLIKYYIDGQK KKCGEERRRVNQFLDYQLQEFGLGVMNT EWIIES with polyhistidine tag at the C-terminus



SDS-PAGE analysis of recombinant human IL-5

产品组成

成分:	从含有 1X PBS, pH 8.0 溶液中冻干的蛋白质.
-----	-------------------------------

产品储存/运输

产品形式	储存温度	储存时间
冻干粉	-20℃至-80℃	自收到之日起1年
重悬液（初始）	2℃至8℃	不超过1周
重悬液（经稀释）	-20℃至-80℃	3到6个月

运输方式： 蓝冰

产品使用

- 1、开盖前，建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉，溶液浓度不低于100μg/mL，不高于1mg/mL，并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液，2-8℃无菌保存不超过1周。
- 4、如需长期保存，推荐使用无菌的含载体蛋白（如0.1% BSA、10% FBS 或5% HSA）的溶液进一步稀释（不低于10ug/mL）后分装保存，-20℃至-80℃无菌保存3到6个月。无血清实验需求时，可更换为5%海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;
FACS= Fluorescence activated Cell Sorting; FA= Functional Assay