

CHAMOT

重组人BMP4拮抗剂 /Recombinant Noggin, Human, Animal-Free

CM172-5HP

CM172-20HP

CM172-100HP

CM172-500HP

CM172-1000HP



CHAMOT

乔默®生物

Specialize In Cytokines

CONTENT

1 产品简介

2 产品储存/运输

3 产品使用

4 实验数据展示

重组人BMP4拮抗剂 /Recombinant Noggin, Human, Animal-Free

产品编号	规格
CM172-5HP	5 ug
CM172-20HP	20 ug
CM172-100HP	100 ug
CM172-500HP	500 ug
CM172-1000HP	1 mg

产品简介**背景描述**

Noggin is a 46.2 kDa bioactive protein, which exists as a disulfide-linked homodimer (each chain 23.1 kDa). Noggin binds members of the transforming Growth Factors-beta (TGF beta) superfamily signaling proteins, such as bone morphogenetic protein-4 (BMP-4), which inactivates their activities. As a extracellular antagonist of BMP proteins, Noggin involves in the development of many body tissues, including nerve tissue, muscles, and bones. In addition, Noggin is able to inhibit chondrocyte differentiation through its interaction with GDF5.

别称

NOG, Noggin, SYM1, symphalangism 1 (proximal), synostoses (multiple) syndrome 1, SYNS1, SYNS1A

蛋白编码

Q13253

分子量

The protein has a calculated MW of 24 kDa. The protein migrates as 27 kDa under reducing condition (SDS-PAGE analysis).

表达系统

Escherichia coli

纯度

>98% as determined by SDS-PAGE.

生物活性

Measure by its ability to inhibit BMP-4-induced alkaline phosphatase production by ATDC5 cells. The ED₅₀ for this effect is <0.05 µg/mL in the presence of 50 ng/mL of recombinant human BMP-4.

内毒素检测

<0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列

MQHYLHIRPAPSDNLPLVDLIEHPDPIFDPKKDLNETLLRSLLGGHYDPGFMATSPPEDRPGGGGGAAGGAEDL
AELDQLLRQRPSGAMPSEIKGLEFSEGLAQGKKQRLSKKLRRKLQMWLWSQTFCPVLYAWNLDLGSRFWPRYV
KVGSCFSKRSCSVPEGMVCKPSKSVHLTVLRWRCQRRGGQRCGWIPYIPISECKCSC with polyhistidine tag at
the C-terminus

产品形式

The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

产品应用

Cell Culture

产品储存/运输

储存

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

运输


Blue Ice

产品使用

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 µg/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

实验数据展示

>



SDS- PAGE analysis of recombinant human Noggin

For Research Use or Further Manufacturing Only

Chamot Biotechnology(Shanghai) Co., Ltd. www.chamot-bio.com

Tel: 021-51880030 Mail: info@chamot-bio.com QQ: 864920491