

# CHAMOT

## Recombinant Human CXCL4

CM162-5HP  
CM162-20HP  
CM162-100HP  
CM162-500HP  
CM162-1000HP



CHAMOT

喬默生物技術(上海)有限公司  
CHAMOT BIOTECHNOLOGY CO., LTD.

# CONTENT

**1** 产品简介

**2** 产品组成

**3** 产品储存

**4** 产品使用


## Recombinant Human CXCL4 (C-X-C motif chemokine4)

编号:	CM162-5HP CM162-20HP CM162-100HP CM162-500HP CM162-1000HP	规格:	5 µg 20 µg 100 µg 500 µg 1 mg
类别:	重组蛋白	应用:	Functional Assay

### 产品简介

**描述:** CXCL4, also known as platelet factor 4 (PF-4), is one of the most plentiful platelet chemokines. Depending on the cell type, CSCL4 may has several biological functions. CXCL4 is mainly produced in megakaryocytes, released from the  $\alpha$ -granules of platelets as a tetramer at micromolar concentrations depending on platelet activation. CXCL4 has both procoagulant and anticoagulant activities, thereby can bind heparin and neutralize the anticoagulant effect of heparin. In addition, CXCL4 also have functions such as inhibiting factor XII, and vitamin K dependent coagulation factor, and stimulating activated protein C generation. As a strong tumor inhibitor, CXCL4 can inhibit endothelial cell migration, proliferation, and in vivo angiogenesis through interfering with the angiogenic effect of growth factors such as FGF and VEGF.

**来源:** *Escherichia coli*  
**纯度:** >98% as determined by SDS-PAGE.



SDS-PAGE analysis of recombinant human CXCL4

生物学活性: Measure by its ability to chemoattract human THP1 cells. Chemotactic activity was observed at a concentration of 10 ng/mL.

内毒素检测: < 0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列: EAEEDGDLQCLCVKTTTSQVRPRHITSLEVIKAGPHCPTAQLIATLKNRGRKICLDLQAPLYKKIIKKLLES with polyhistidine tag at the N-terminus

## 产品组成

成分: 从含有 1X PBS, pH 7.4 溶液中冻干的蛋白质.

## 产品储存/运输

产品形式	储存温度	储存时间
冻干粉	-20°C至-80°C	自收到之日起1年
重悬液 (初始)	2°C至8°C	不超过1周
重悬液 (经稀释)	-20°C至-80°C	3到6个月

运输方式: 蓝冰

## 产品使用

- 1、开盖前, 建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉, 溶液浓度不低于100µg/mL, 不高于1mg/mL, 并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液, 2 -8°C无菌保存不超过1周。
- 4、如需长期保存, 推荐使用无菌的含载体蛋白 (如0.1 %BSA、10 %FBS 或5 %HSA ) 的溶液进一步稀释 ( 不低于10ug/mL ) 后分装保存, -20°C至 -80°C无菌保存3到6个月。无血清实验需求时, 可更换为 5%海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;  
FACS= Fluorescence activated Cell Sorting; FA= Functional Assay