

CHAMOT

Recombinant Human IL-17A

CM018-5HP
CM018-20HP
CM018-100HP
CM018-500HP
CM018-1000HP



CHAMOT

喬默生物技術(上海)有限公司
CHAMOT BIOTECHNOLOGY CO., LTD.

CONTENT

1 产品简介

2 产品组成

3 产品储存

4 产品使用

Recombinant Human IL-17A (Interleukin-17A)

编号:	CM018-5HP CM018-20HP CM018-100HP CM018-500HP CM018-1000HP	规格:	5 µg 20 µg 100 µg 500 µg 1 mg
类别:	重组蛋白	应用:	Functional Assay

产品简介

描述: Interleukin-17A is a protein that in humans is encoded by the IL17A gene. The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO).


来源: *Escherichia coli*

纯度: >98% as determined by SDS-PAGE. Ni-NTA chromatography

生物学活性: Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED₅₀ for this effect is < 6 ng/mL.

内毒素检测: < 0.01 EU per 1 µg of the protein by the LAL method.

蛋白序列: MIVKAGITIPRNP GCPNSEDKNFPRTV
MVNLNIHNRNTNTNPKRSSDYNNRS
TSPWNLHRNEDPERYPSVIWEAKCR
HLGCINADGNVDYHMNSVPIQQEIL
VLRREPPHCPNSFRLEKILVSVGCTCV
TPIVHHVA with polyhistidine tag at the C-terminus



SDS-PAGE analysis of recombinant human IL-17A

产品组成

成分: 从含有 1X PBS, pH 8.0 溶液中冻干的蛋白质.

产品储存

产品形式	储存温度	储存时间
冻干粉	-20℃至-80℃	自收到之日起1年
重悬液（初始）	2℃至8℃	不超过1周
重悬液（经稀释）	-20℃至-80℃	3到6个月

运输方式： 蓝冰

产品使用

- 1、开盖前，建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉，溶液浓度不低于100μg/mL，不高于1mg/mL，并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液，2-8℃无菌保存不超过1周。
- 4、如需长期保存，推荐使用无菌的含载体蛋白（如0.1% BSA、10% FBS 或5% HSA）的溶液进一步稀释（不低于10ug/mL）后分装保存，-20℃至-80℃无菌保存3到6个月。无血清实验需求时，可更换为5%海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;
FACS= Fluorescence activated Cell Sorting; FA= Functional Assay