

CHAMOT

Recombinant Human IL-8

CM010-5HP
CM010-20HP
CM010-100HP
CM010-500HP
CM010-1000HP



CHAMOT

喬默生物技術(上海)有限公司
CHAMOT BIOTECHNOLOGY CO. LTD.

CONTENT

1 产品简介

2 产品组成

3 产品储存

4 产品使用

Recombinant Human IL-8 (aa 28-99) (Interleukin-8)

编号:	CM010-5HP	规格:	5 µg
	CM010-20HP		20 µg
	CM010-100HP		100 µg
	CM010-500HP		500 µg
	CM010-1000HP		1 mg

类别: 重组蛋白 应用: Functional Assay

产品简介

描述: Interleukin 8 (IL8 or chemokine (C-X-C motif) ligand 8, CXCL8) is a chemokine produced by macrophages and other cell types such as epithelial cells, airway smooth muscle cells, and endothelial cells. Endothelial cells store IL-8 in their storage vesicles, the Weibel-Palade bodies. In humans, the interleukin-8 protein is encoded by the CXCL8 gene. IL-8 is initially produced as a precursor peptide of 99 amino acids which then undergoes cleavage to create several active IL-8 isoforms. In culture, a 72 amino acid peptide is the major form secreted by macrophages.

来源:

Escherichia coli

纯度:

>98% as determined by SDS-PAGE.

生物学活性:


Measure by its ability to chemoattract PBMC using a concentration range of 1-10 ng/mL.

内毒素检测:

< 0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列:

MSAKELRCQCICIKTYSKPFHPKFIKELRV
IESGPHCANTEIIVKLSDGRELCLDPKE
NWVQRVVEKFLKRAENS with polyhistidine tag at the C-terminus



SDS-PAGE analysis of recombinant human IL-8

产品组成

成分：从含有 1X PBS, pH 8.0 溶液中冻干的蛋白质.

产品储存/运输

产品形式	储存温度	储存时间
冻干粉	-20°C至-80°C	自收到之日起1年
重悬液 (初始)	2°C至8°C	不超过1周
重悬液 (经稀释)	-20°C至-80°C	3到6个月

运输方式： 蓝冰

产品使用

- 1、开盖前，建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉，溶液浓度不低于100μg/mL，不高于1mg/mL，并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液，2 -8°C无菌保存不超过1周。
- 4、如需长期保存，推荐使用无菌的含载体蛋白（如0.1% BSA、10% FBS 或 5% HSA）的溶液进一步稀释（不低于10ug/mL）后分装保存，-20°C至 -80°C无菌保存3到6个月。无血清实验需求时，可更换为 5% 海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;
 FACS= Fluorescence activated Cell Sorting; FA= Functional Assay