

# CHAMOT

## Recombinant Mouse IL-17B

CM053-5MP  
CM053-20MP  
CM053-100MP  
CM053-500MP  
CM053-1000MP



CHAMOT

喬默生物技術(上海)有限公司  
CHAMOT BIOTECHNOLOGY CO., LTD.

# CONTENT

1 产品简介

2 产品组成

3 产品储存

4 产品使用

## Recombinant Mouse IL-17B (Interleukin-17B)

编号:	CM053-5MP	规格:	5 µg
	CM053-20MP		20 µg
	CM053-100MP		100 µg
	CM053-500MP		500 µg
	CM053-1000MP		1 mg
类别:	重组蛋白	应用:	Functional Assay

### 产品简介

**描述:** The interleukin (IL)-17 superfamily, a relatively new family of cytokines, consists of six ligands (from IL-17A to IL-17F), which bind to five receptor subtypes (from IL-17RA to IL-17RE) and induce downstream signaling.  
**L17B Diseases associated with IL17B** include spondyloarthropathy and neuronitis, and among its related super-pathways are Mucin expression in CF via IL-6, IL-17 signaling pathways, and STAT3 Pathway.


**来源:** *Escherichia coli*

**纯度:** >98% as determined by SDS-PAGE.  
 Ni-NTA chromatography.

**生物学活性:** Measure by its ability to induce IL-8 secretion in HepG2 cells. The ED<sub>50</sub> for this effect is < 1.5 ng/mL.

**内毒素检测:** < 0.1 EU per 1 µg of the protein by the LAL method.

**蛋白序列:** MHPRNTKGKRKGQGRPSPLAPGPH  
 QVPLDLVSRVKPYARMEYERNLGEM  
 VAQLRNSSEPAKKCEVNQLQLWLSN  
 KRSLSPWGYSINHDPSRIPADLPEARC  
 LCLGCVNPFTMQEDRSMVSVPVFSQ  
 VPVRRRLCPQPPRPGPCRQRVVMETI  
 AVGCTCIF with polyhistidine tag at the C-terminus



SDS-PAGE analysis of recombinant mouse IL-17B

## 产品组成

成分： 从含有 1X PBS, pH 7.4 溶液中冻干的蛋白质.

## 产品储存/运输

产品形式	储存温度	储存时间
冻干粉	-20°C至-80°C	自收到之日起1年
重悬液 (初始)	2°C至8°C	不超过1周
重悬液 (经稀释)	-20°C至-80°C	3到6个月

运输方式： 蓝冰

## 产品使用

- 1、开盖前，建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉，溶液浓度不低于100μg/mL，不高于1mg/mL，并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液，2-8°C无菌保存不超过1周。
- 4、如需长期保存，推荐使用无菌的含载体蛋白（如0.1% BSA、10% FBS 或 5% HSA）的溶液进一步稀释（不低于10ug/mL）后分装保存，-20°C至 -80°C无菌保存3到6个月。无血清实验需求时，可更换为 5% 海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;  
 FACS= Fluorescence activated Cell Sorting; FA= Functional Assay