

CHAMOT

Recombinant Mouse IL-22

CM019-5MP
CM019-20MP
CM019-100MP
CM019-500MP
CM019-1000MP



CHAMOT

喬默生物技術(上海)有限公司
CHAMOT BIOTECHNOLOGY CO., LTD.

CONTENT

1 产品简介

2 产品组成

3 产品储存

4 产品使用

Recombinant Mouse IL-22 (Interleukin-22)

编号:	CM019-5MP	规格:	5 µg
	CM019-20MP		20 µg
	CM019-100MP		100 µg
	CM019-500MP		500 µg
	CM019-1000MP		1 mg

类别: 重组蛋白 应用: Functional Assay

产品简介


描述: Interleukin-22 (IL-22) a member of a group of cytokines called the IL-10 family or IL-10 superfamily (including IL-19, IL-20, IL-24, and IL-26), a class of potent mediators of cellular inflammatory responses. It shares use of IL-10R2 in cell signaling with other members of this family, IL-10, IL-26, IL-28A/B and IL-29. IL-22 is produced by activated NK and T cells and initiates innate immune responses against bacterial pathogens especially in epithelial cells such as respiratory and gut epithelial cells. IL-22 along with IL-17 is rapidly produced by splenic LTi-like cells and also produced by Th17 cells and likely plays a role in the coordinated response of both adaptive innate immune systems, autoimmunity and tissue regeneration.

来源: *Escherichia coli*

纯度: >98% as determined by SDS-PAGE.
Ni-NTA chromatography.

生物学活性: Measure by its ability to induce IL-10 secretion in COLO205 cells. The ED₅₀ for this effect is < 0.3 ng/mL.

内毒素检测: < 0.1 EU per 1 µg of the protein by the LAL method.



SDS-PAGE analysis of recombinant mouse IL-22

蛋白序列：
 MLPVNTRCKLEVSNFQQPYIVNRTFM
 LAKEASLADNNNTDVRLIGEKLFRGVSA
 KDQCYLMKQVLNFTLEDVLLPQSDF
 QPYMQEVVPFLTKLSNQLSSCHISGD
 DQNIQKNVRRLKETVKKLGESGEIKAI
 GELDLLFMSLRNACV with
 polyhistidine tag at the C-terminus

产品组成

成分：从含有 1X PBS, pH 7.4 溶液中冻干的蛋白质.

产品储存/运输

产品形式	储存温度	储存时间
冻干粉	-20°C至-80°C	自收到之日起1年
重悬液 (初始)	2°C至8°C	不超过1周
重悬液 (经稀释)	-20°C至-80°C	3到6个月

运输方式： 蓝冰

产品使用

- 1、开盖前，建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉，溶液浓度不低于100μg /mL，不高于1mg/mL，并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液，2-8°C无菌保存不超过1周。
- 4、如需长期保存，推荐使用无菌的含载体蛋白（如0.1 %BSA、10% FBS 或 5% HSA）的溶液进一步稀释（不低于10ug/mL）后分装保存，-20°C至 -80°C无菌保存3到6个月。无血清实验需求时，可更换为 5% 海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;
 FACS= Fluorescence activated Cell Sorting; FA= Functional Assay