

# CHAMOT

## Recombinant Mouse IL-33

CM030-5MP  
CM030-20MP  
CM030-100MP  
CM030-500MP  
CM030-1000MP



CHAMOT

喬默生物技術(上海)有限公司  
CHAMOT BIOTECHNOLOGY CO., LTD.

# CONTENT

1 产品简介

2 产品组成

3 产品储存

4 产品使用

## Recombinant Mouse IL-33 (Interleukin-33)

编号:	CM030-5MP CM030-20MP CM030-100MP CM030-500MP CM030-1000MP	规格:	5 µg 20 µg 100 µg 500 µg 1 mg
-----	---	-----	---

类别: 重组蛋白 应用: Functional Assay

### 产品简介

描述: Interleukin 33 (IL-33) is a member of the IL-1 family that potently drives production of T helper-2 (Th2)-associated cytokines (e.g., IL-4). IL33 is a ligand for ST2 (IL1RL1), an IL-1 family receptor that is highly expressed on Th2 cells, mast cells and group 2 innate lymphocytes. It is expressed by a wide variety of cell types, including fibroblasts, mast cells, dendritic cells, macrophages, osteoblasts, endothelial cells, and epithelial cells.


来源: *Escherichia coli*

纯度: >98% as determined by SDS-PAGE.  
Ni-NTA chromatography.

生物学活性: Measure by its ability to induce proliferation in D10.G4.1 cells. The ED<sub>50</sub> for this effect is < 40 pg/mL. The specific activity of recombinant mouse IL-33 is > 2 x 10<sup>7</sup> IU/mg.

内毒素检测: < 0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列: MSIQGTSLLTQSPASLSTYNDQSVSFV  
LENGCYVINVDDSGKDQECDQVLLR  
YYESPCPASQSGDGVDGKKLMVNMS  
PIKDTDIWLHANDKDYSVELQRGDVS  
PPEQAFFVLHKKSSDFVSFECKNLPGT  
YIGVKDNQLALVEEKDESCNNIMFKLS  
KI with polyhistidine tag at the C-terminus



SDS-PAGE analysis of recombinant mouse IL-33

## 产品组成

成分： 从含有 1X PBS, pH 7.4 溶液中冻干的蛋白质.

## 产品储存/运输

产品形式	储存温度	储存时间
冻干粉	-20°C至-80°C	自收到之日起1年
重悬液 (初始)	2°C至8°C	不超过1周
重悬液 (经稀释)	-20°C至-80°C	3到6个月

运输方式： 蓝冰

## 产品使用

- 1、开盖前，建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉，溶液浓度不低于100μg/mL，不高于1mg/mL，并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液，2-8°C无菌保存不超过1周。
- 4、如需长期保存，推荐使用无菌的含载体蛋白（如0.1%BSA、10%FBS 或5%HSA）的溶液进一步稀释（不低于10ug/mL）后分装保存，-20°C至-80°C无菌保存3到6个月。无血清实验需求时，可更换为 5%海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;  
 FACS= Fluorescence activated Cell Sorting; FA= Functional Assay