

# CHAMOT

## Recombinant Mouse TNF alpha

CM038-5MP

CM038-20MP

CM038-100MP

CM038-500MP

CM038-1000MP



CHAMOT

喬默生物技術(上海)有限公司  
CHAMOT BIOTECHNOLOGY CO., LTD.

# CONTENT

1 产品简介

2 产品组成

3 产品储存

4 产品使用

## Recombinant Mouse TNF alpha (Tumor necrosis factor alpha)

编号:	CM038-5MP	规格:	5 µg
	CM038-20MP		20 µg
	CM038-100MP		100 µg
	CM038-500MP		500 µg
	CM038-1000MP		1 mg

类别: 重组蛋白 应用: Functional Assay

### 产品简介

描述: TNF- $\alpha$  is a kind of pleiotropic pro-inflammatory cytokine. It is secreted by various cells, such as adipocytes, activated monocytes, macrophages, B cells, T cells and fibroblasts. Proteolysis of the integral membrane precursor form of TNF- $\alpha$  from cells soluble can release homotrimeric TNF- $\alpha$ . TNF- $\alpha$  can bind with some TNF- $\alpha$  receptors induces apoptosis, besides, also trigger other responses depending on cell type, receptor expression, and signal transduction status. TNF- $\alpha$  participate in the inflammatory response.


来源: *Escherichia coli*

纯度: >98% as determined by SDS-PAGE.  
Ni-NTA chromatography.

生物学活性: Measure by its ability to induce cytotoxicity in L929 cells in the presence of actinomycin D. The ED<sub>50</sub> for this effect is < 40 pg/mL. The specific activity of recombinant mouse TNF alpha is approximately >2.5x 10<sup>7</sup> IU/mg.

内毒素检测: < 0.1 EU per 1 µg of the protein by the LAL method.

蛋白序列: MLRSSSQNSSDKPVAHVANHQVEE  
QLEWLSQRANALLANGMDLKDNLQL  
VVPADGLYLVYSQVLFKGQGCPDYVL  
LTHTVSRAISYQEKVNLLSAVKSPCP  
KDTPEGAEELKPWYEPIYLGGVFQLEGD



SDS-PAGE analysis of recombinant mouse TNF alpha

QLSAEVNLPKYLDFAESGQVYFGVIAL  
 with polyhistidine tag at the C-  
 terminus

交叉反应种属: Rat(94.27%)

## 产品组成

成分: 从含有 1X PBS, pH 8.0 溶液中冻干的  
 蛋白质.

## 产品储存/运输

产品形式	储存温度	储存时间
冻干粉	-20°C至-80°C	自收到之日起1年
重悬液 (初始)	2°C至8°C	不超过1周
重悬液 (经稀释)	-20°C至-80°C	3到6个月

运输方式: 蓝冰

## 产品使用

- 1、开盖前，建议3000-3500rpm离心5min。
- 2、推荐使用无菌水重悬冻干粉，溶液浓度不低于100μg/mL，不高于1mg/mL，并室温静置至少20min以充分溶解。勿涡旋剧烈振荡。
- 3、重悬后的溶液，2-8°C无菌保存不超过1周。
- 4、如需长期保存，推荐使用无菌的含载体蛋白（如0.1%BSA、10%FBS 或 5%HSA）的溶液进一步稀释(不低于10ug/mL)后分装保存，-20°C至-80°C无菌保存3到6个月。无血清实验需求时，可更换为 5%海藻糖溶液作为载体。避免反复冻融。

WB= Western Blot; IP= Immunoprecipitation; IF= Immunofluorescence; IHC= Immunohistochemistry;  
 FACS= Fluorescence activated Cell Sorting; FA= Functional Assay